

# NEW RESULTS FOR THE LIEBAU PHENOMENON VIA FIXED POINT INDEX

J. A. CID, G. INFANTE, M. TVRDÝ, AND M. ZIMA

ABSTRACT. We prove new results regarding the existence of positive solutions for a nonlinear periodic boundary value problem related to the Liebau phenomenon. As a consequence we obtain new sufficient conditions for the existence of a pump in a simple model. Our methodology relies on the use of classical fixed point index. Some examples are provided to illustrate our theory. We improve and complement previous results in the literature.

## 1. INTRODUCTION

During the experiments developed in the 1950s, the German cardiologist Gerhart Liebau observed (see [1]) that a periodic compression could produce the circulation of a fluid in a mechanical system without valves to ensure the direction of the flow. This valveless pumping effect is nowadays called the Liebau phenomenon. It was reported to occur for instance in embryonic blood circulation, in applications of nanotechnology and in oceanic currents, see e.g. [2, 3, 4, 5, 6] or [7, Chapter 8]. In particular, G. Propst [6] presented an explanation of the pumping effect for flow configurations of several rigid tanks that are connected by rigid pipes. He proved the existence of periodic solutions to the corresponding differential equations for systems of 2 or 3 tanks. However, the apparently simplest configuration consisting of 1 pipe and 1 tank turned out to be, from mathematical point of view, the most interesting one, as it leads to the singular periodic problem

$$(1.1) \quad \begin{cases} u''(t) + a u'(t) = \frac{1}{u(t)} (e(t) - b(u'(t))^2) - c, & t \in [0, T], \\ u(0) = u(T), \quad u'(0) = u'(T), \end{cases}$$

where  $u'$  is the fluid velocity in the pipe (oriented in the direction from the tank to the piston),  $T > 0$ ,

$$a = \frac{r_0}{\rho}, \quad b = 1 + \frac{\zeta}{2}, \quad c = \frac{gA_\pi}{A_\tau}, \quad e(t) = \frac{gV_0}{A_\tau} - \frac{p(t)}{\rho},$$

---

2010 *Mathematics Subject Classification.* Primary 34B18, secondary 34B27, 34B60.

*Key words and phrases.* Valveless pumping; Periodic boundary value problem; Cone; Fixed point index; Green's function.

$r_0$  is the friction coefficient,  $\rho$  is the density of the fluid,  $\zeta \geq 1$  is the junction coefficient (depending on the particular geometry and smoothness of the junction of the tank and the pipe),  $g$  is the acceleration of gravity,  $A_\tau$  is the cross section of the tank,  $A_\pi$  is the cross section of the pipe (small in comparison with  $A_\tau$ ),  $V_0$  is the total volume (assumed to be constant) of the fluid in the system and  $p$  is the  $T$ -periodic external force. As a result, from the fluid mechanics point of view, the assumptions

$$a \geq 0, \quad b > 1, \quad c > 0, \quad e \text{ continuous and } T - \text{periodic}$$

are quite natural, from  $\zeta \geq 1$  we would even have  $b \geq 3/2$ . Of course, we are interested in the search of positive solutions of problem (1.1). A detailed justification of the model can be also found e.g. in [7, Chapter 8].

One can observe that if a periodic external force  $e$  produces a nonconstant periodic response  $u$  then the mean level of the fluid in the tank is higher than the level produced by a constant force with the same mean value. Moreover the increasing of the level is proportional to  $\|u'\|^2$ .

The change of variables  $u = x^\mu$ , where  $\mu = \frac{1}{b+1}$ , was used in [8] in order to overcome the singularity, transforming and simplifying problem (1.1) into the regular BVP

$$(1.2) \quad \begin{cases} x''(t) + a x'(t) = \frac{e(t)}{\mu} x^{1-2\mu}(t) - \frac{c}{\mu} x^{1-\mu}(t), & t \in [0, T], \\ x(0) = x(T), \quad x'(0) = x'(T), \end{cases}$$

where  $0 < \mu < \frac{1}{2}$ . By means of the lower and upper solution technique Cid and co-authors [8] provided results on the existence and stability of a positive solution of (1.2).

In our recent paper [9] we considered a generalization of problem (1.2), namely

$$(1.3) \quad \begin{cases} x''(t) + ax'(t) = r(t) x^\alpha(t) - s(t) x^\beta(t), & t \in [0, T], \\ x(0) = x(T), \quad x'(0) = x'(T), \end{cases}$$

under the assumption

$$(H0) \quad a \geq 0, \quad r, s : [0, T] \rightarrow \mathbb{R} \text{ are continuous and } 0 < \alpha < \beta < 1.$$

Of course, to extend the obtained solution of the boundary value problem (1.3) (on  $[0, T]$ ) to a  $T$ -periodic solution of the corresponding differential equation, we would have to assume that  $r$  and  $s$  are also  $T$ -periodic. Making use of a shifting argument and Krasnosel'skiĭ's expansion/compression fixed point theorem on cone, we succeeded in [9] to improve the existence results from [8].

Furthermore, Torres in [7, Chapter 8] obtained a priori bounds for the periodic solutions of (1.1), which together with the Brouwer degree theoretical arguments, led to an alternative existence result.

We point out that the assumption

$$\min_{t \in [0, T]} e(t) > 0$$

is a common feature of the existence results in [7, 8, 9]. The goal of this paper is twofold: first, to improve the main results from [9] and, second, to obtain explicit sufficient conditions for the existence of periodic solutions of problem (1.2) that allow, for the first time, the function  $e$  to be sign-changing. Similarly to the papers [10, 11, 12, 13, 14, 15], our main tool will be the classical fixed point index on cones.

The paper is organized as follows: in Section 2 we present the shifting argument and recall some known facts regarding the Green's function of the problem and some properties of the fixed point index. In Section 3 we perform the fixed point index calculations that we use to prove our main results. In Section 4 we present the main results, some consequences and illustrative examples.

## 2. PRELIMINARIES

By means of a shifting argument (see [9]) problem (1.3) may be rewritten in the equivalent form

$$(2.1) \quad \begin{cases} x''(t) + a x'(t) + m^2 x(t) = r(t) x^\alpha(t) - s(t) x^\beta(t) + m^2 x(t), & t \in [0, T], \\ x(0) = x(T), \quad x'(0) = x'(T), \end{cases}$$

with  $m \in \mathbb{R}$ . In the sequel, we denote the right-hand side of the differential equation in (2.1) by  $f_m$ , i.e.

$$(2.2) \quad f_m(t, x) = r(t) x^\alpha - s(t) x^\beta + m^2 x \quad \text{for } t \in [0, T] \text{ and } x \in [0, +\infty).$$

We will assume that the linearization

$$(2.3) \quad \begin{cases} x''(t) + a x'(t) + m^2 x(t) = h(t), & t \in [0, T], \\ x(0) = x(T), \quad x'(0) = x'(T), \end{cases}$$

of (2.1) possesses a positive Green's function. Its existence and its further needed properties are given by the following lemma.

**Lemma 1.** *Assume that*

$$(2.4) \quad a \geq 0 \quad \text{and} \quad 0 < m < \sqrt{\left(\frac{\pi}{T}\right)^2 + \left(\frac{a}{2}\right)^2}.$$

*Then there exists a unique function  $G_m(t, s)$  defined on  $[0, T] \times [0, T]$  such that*

$$(G1) \quad \left\{ \begin{array}{l} \text{For each } h \text{ continuous on } [0, T], \text{ the function} \\ x(t) = \int_0^T G_m(t, s) h(s) \, ds \quad \text{for } t \in [0, T] \\ \text{is the unique solution of (2.3).} \end{array} \right.$$

$$(G2) \quad G_m > 0 \text{ on } [0, T] \times [0, T].$$

$$(G3) \quad \int_0^T G_m(t, s) \, ds = \frac{1}{m^2} \quad \text{for all } t \in [0, T].$$

$$(G4) \quad \left\{ \begin{array}{l} \text{There exists a constant } c_m \in (0, 1) \text{ such that} \\ G_m(t, s) \geq G_m(s, s) \geq c_m G_m(t, s) \quad \text{for all } (t, s) \in [0, T] \times [0, T]. \end{array} \right.$$

*Proof.* The existence of a function  $G_m$  with the properties (G1) and (G2) is given by [16, Proposition 2.2] and for the rest, see [9, Appendix A].  $\square$

Throughout, given  $T$ ,  $a$  and  $m$  in (2.4), we will write  $G_m$  and  $c_m$  for the corresponding function  $G_m$  in (G1) and the corresponding constant  $c_m$  in (G4).

Furthermore, let us recall that a *cone*  $P$  in a Banach space  $X$  is a closed, convex subset of  $X$  such that  $P \cap (-P) = \{0\}$  and  $\lambda x \in P$  for  $x \in P$  and  $\lambda \geq 0$ . Here we will work in the space  $X = \mathcal{C}[0, T]$  of continuous functions on  $[0, T]$  endowed with the usual maximum norm  $\|x\| = \max\{|x(t)| : t \in [0, T]\}$ . The properties of Green's function  $G_m$  of (2.3) stated in Lemma 1 enable us to use the cone

$$(2.5) \quad P = \{x \in X : x(t) \geq c_m \|x\| \text{ for } t \in [0, T]\},$$

a type of a cone first used by M.A. Krasnosel'skiĭ and D. Guo, see for example [17, 18].

Whenever (2.4) is true, we define the operator  $F : P \rightarrow X$  by

$$(2.6) \quad Fx(t) = \int_0^T G_m(t, s) f_m(s, x(s)) \, ds \quad \text{for } x \in P \text{ and } t \in [0, T],$$

where  $G_m$  is the corresponding Green's function for (2.3). Then, any fixed point of  $F$  in  $P$  is a nonnegative solution of problem (2.1) and, simultaneously, of problem (1.3). In order

to obtain the existence of a fixed point of this kind we make use of the classical fixed point index. For readers' convenience we formulate below the principles that we will need later.

If  $\Omega$  is an open bounded subset of  $P$  (in the relative topology) we denote by  $\overline{\Omega}$  and  $\partial\Omega$  the closure and the boundary relative to  $P$ . When  $\Omega$  is an open bounded subset of  $X$  we write  $\Omega_P = \Omega \cap P$ . Recall that for a given cone  $P$ , an open set  $\Omega$  and a compact operator  $F: \overline{\Omega_P} \rightarrow P$ , the symbol  $i_P(F, \Omega_P)$  stands for the fixed point index of  $F$  with respect to  $P$  and  $\Omega$ . For the definition and more details concerning the properties of the fixed point index see e.g. [18, 19, 20, 21]. The following existence principle is well-known, but we include its short proof for completeness.

**Theorem 2.** *Let  $P$  be a cone in a Banach space  $X$ . Let  $\Omega, \Omega' \subset X$  be open bounded sets such that  $0 \in \Omega'_P$  and  $\overline{\Omega'_P} \subset \Omega_P$ . Assume that  $F: \overline{\Omega_P} \rightarrow P$  is a compact map such that  $Fx \neq x$  for  $x \in \partial\Omega_P \cup \partial\Omega'_P$ ,  $i_P(F, \Omega_P) = 1$  and  $i_P(F, \Omega'_P) = 0$ . Then  $F$  has a fixed point in  $\Omega_P \setminus \overline{\Omega'_P}$ .*

*Proof* It follows from the additivity and solution properties of the fixed point index, cf. e.g. [18, Theorem 2.3.1 and Theorem 2.3.2]. For an analogous argument, see also e.g. the proof of Theorem 12.3 in [19].  $\square$

To verify the assumptions of Theorem 2, the following assertion will be helpful.

**Theorem 3** ([18], Lemma 2.3.1 and Corollary 2.3.1). *Let  $P$  be a cone in a Banach space  $X$ ,  $\Omega$  be an open bounded set such that  $0 \in \Omega_P$  and let  $F: \overline{\Omega_P} \rightarrow P$  be compact. Then*

- (i) *If  $Fx \neq \lambda x$  for all  $x \in \partial\Omega_P$  and all  $\lambda \geq 1$ , then  $i_P(F, \Omega_P) = 1$ .*
- (ii) *If there exists  $x_0 \in P \setminus \{0\}$  such that  $x - Fx \neq \lambda x_0$  for all  $x \in \partial\Omega_P$  and all  $\lambda \geq 0$ , then  $i_P(F, \Omega_P) = 0$ .*

We will complete this section by introducing further notations needed later:

For a given continuous function  $h: [0, T] \rightarrow \mathbb{R}$  we denote

$$\bar{h} = \frac{1}{T} \int_0^T h(s) \, ds, \quad h_* = \min\{h(t) : t \in [0, T]\}, \quad h^* = \max\{h(t) : t \in [0, T]\},$$

and

$$h_+(t) = \max\{h(t), 0\} \quad \text{for } t \in [0, T].$$

## 3. CALCULATIONS OF THE FIXED POINT INDEX

We will assume that (H0) holds and

$$(H1) \quad \begin{cases} \text{there are } m > 0, R_2 > 0 \text{ and } R_1 \in (0, R_2) \text{ such that} \\ m^2 < \left(\frac{\pi}{T}\right)^2 + \left(\frac{a}{2}\right)^2 \text{ and } f_m(t, x) \geq 0 \text{ for } t \in [0, T] \text{ and } x \in [c_m R_1, R_2], \end{cases}$$

where  $f_m$  is given by (2.2). Let  $\tilde{f}_m: [0, T] \times [0, R_2] \rightarrow \mathbb{R}$  be an arbitrary continuous and nonnegative function which coincides with  $f_m$  on  $[0, T] \times [c_m R_1, R_2]$ .

For  $\rho > 0$  and  $P$  as in (2.5), we define

$$B_\rho = \{x \in P : \|x\| < \rho\} \quad \text{and} \quad B'_\rho = \{x \in P : x_* < c_m \rho\}.$$

Let us note that the sets of the form  $B'_\rho$  were introduced and utilized already by Lan in [11].

Note that we have  $B'_\rho \subset B_\rho$ .

We will consider the operator

$$(3.1) \quad \tilde{F}x(t) = \int_0^T G_m(t, s) \tilde{f}_m(s, x(s)) \, ds \quad \text{for } x \in \overline{B_{R_2}}.$$

where  $G_m$  is Green's function of (2.3) whose existence and properties are given by Lemma 1.

The main object of this Section is to yield sufficient conditions for the fixed point index of  $\tilde{F}$  to be 1 or 0. In order to do this, we utilize, in the spirit of the papers by Lan [12] and Webb [15], the explicit dependence of the nonlinearity  $f_m$  on  $t$ .

**Lemma 4.** *Assume (H0) and (H1). Then the operator  $\tilde{F}$  maps  $\overline{B_{R_2}}$  into  $P$  and is compact.*

*Proof.* For  $x \in \overline{B_{R_2}}$  and  $t \in [0, T]$ , we have by Lemma 1 and construction of  $\tilde{f}_m$

$$\tilde{F}x(t) = \int_0^T G_m(t, s) \tilde{f}_m(s, x(s)) \, ds \geq \int_0^T G_m(s, s) \tilde{f}_m(s, x(s)) \, ds$$

and

$$\int_0^T G_m(s, s) \tilde{f}_m(s, x(s)) \, ds \geq c_m \int_0^T G_m(t, s) \tilde{f}_m(s, x(s)) \, ds.$$

Therefore  $\tilde{F}x(t) \geq c_m \|\tilde{F}x\|$  on  $[0, T]$ . Hence  $\tilde{F}(\overline{B_{R_2}}) \subset P$ . The compactness of  $\tilde{F}$  follows in a standard way from the Arzelà-Ascoli theorem.  $\square$

**Lemma 5.** *Assume that (H0) and (H1) hold,  $\tilde{F}x \neq x$  for  $x \in \partial B_{R_2}$  and there exists a continuous function  $g_1$  such that*

$$(H2) \quad f_m(t, x) \leq g_1(t) \quad \text{for } t \in [0, T] \text{ and } x \in [c_m R_2, R_2]$$

and moreover either

$$(H3) \quad \delta_* \leq c_m R_2,$$

or

$$(H4) \quad \delta^* \leq R_2,$$

where

$$\delta(t) = \int_0^T G_m(t, s) g_1(s) ds \quad \text{for } t \in [0, T].$$

Then  $i_P(\tilde{F}, B_{R_2}) = 1$ .

*Proof.* It follows from Lemma 4 that  $\tilde{F}$  maps  $\overline{B_{R_2}}$  into  $P$  and is compact. We show that  $\tilde{F}x \neq \lambda x$  for  $x \in \partial B_{R_2}$  and  $\lambda \geq 1$ , which by Theorem 3 (i) (where we put  $\Omega_P = B_{R_2}$ ) yields  $i_P(\tilde{F}, B_{R_2}) = 1$ .

If not, there exist  $\lambda \geq 1$  and  $x \in P$  with  $\|x\| = R_2$  such that  $\tilde{F}x = \lambda x$ . Since we suppose that  $\tilde{F}x \neq x$  for  $x \in \partial B_{R_2}$ , we may assume that  $\lambda > 1$ . Consequently,  $c_m R_2 \leq x(t) \leq R_2$  for  $t \in [0, T]$ , and, since  $G_m(t, s) > 0$ , hypothesis (H2) implies

$$\lambda x(t) = \int_0^T G_m(t, s) \tilde{f}_m(s, x(s)) ds \leq \int_0^T G_m(t, s) g_1(s) ds = \delta(t).$$

If (H3) holds then we obtain

$$\lambda c_m R_2 \leq \lambda x_* = \min_{t \in [0, T]} \int_0^T G_m(t, s) \tilde{f}_m(s, x(s)) ds \leq \delta_* \leq c_m R_2,$$

a contradiction since  $\lambda > 1$ . If (H4) holds, then taking the maximum on  $[0, T]$  we obtain  $\lambda R_2 = \lambda x^* \leq \delta^* \leq R_2$ , a contradiction.  $\square$

**Lemma 6.** *Assume that (H0) and (H1) hold,  $\tilde{F}x \neq x$  for  $x \in \partial B'_{R_1}$  and there exists a continuous function  $g_0$  such that*

$$(H5) \quad f_m(t, x) \geq g_0(t) \geq 0 \quad \text{for } t \in [0, T] \quad \text{and } x \in [c_m R_1, R_1],$$

and moreover either

$$(H6) \quad \gamma_* \geq c_m R_1,$$

or

$$(H7) \quad \gamma^* \geq R_1,$$

where

$$\gamma(t) = \int_0^T G_m(t, s) g_0(s) ds \quad \text{for } t \in [0, T].$$

Then  $i_P(\tilde{F}, B'_{R_1}) = 0$ .

*Proof.* By Lemma 4, the operator  $\tilde{F} : \overline{B'_{R_1}} \subset \overline{B_{R_1}} \rightarrow P$  is compact. Put  $x_0(t) = 1$  for  $t \in [0, T]$ . Then  $x_0 \in P$  and we shall prove that  $x - \tilde{F}x \neq \lambda x_0$  for all  $x \in \partial B'_{R_1}$  and  $\lambda \geq 0$ , which by Theorem 3 (ii) (where we put  $\Omega'_P = B'_{R_1}$ ) implies that  $i_P(\tilde{F}, B'_{R_1}) = 0$ .

If not, there exist  $x \in \partial B'_{R_1}$  and  $\lambda \geq 0$  such that  $x = \tilde{F}x + \lambda x_0$ . Due to our assumption  $\tilde{F}x \neq x$  for  $x \in \partial B'_{R_1}$ , it is enough to consider  $\lambda > 0$ . Furthermore, note that  $x \in \partial B'_{R_1}$  means that  $x \in P$  and  $x_* = c_m R_1$ .

In particular,  $x \in \partial B'_{R_1}$  implies that  $c_m R_1 \leq x(t) \leq R_1$  for  $t \in [0, T]$ . Then due to (H5) and the positivity of Green's function  $G_m$  we have for  $t \in [0, T]$  that

$$\begin{aligned} x(t) &= \tilde{F}x(t) + \lambda x_0(t) = \int_0^T G_m(t, s) \tilde{f}_m(s, x(s)) ds + \lambda \\ &\geq \int_0^T G_m(t, s) g_0(s) ds + \lambda = \gamma(t) + \lambda. \end{aligned}$$

Now, if (H6) holds then, using Lemma 1 we get the contradiction

$$c_m R_1 = x_* \geq \gamma_* + \lambda > \gamma_* \geq c_m R_1.$$

On the other hand, if (H7) holds, using again Lemma 1 we arrive at the contradiction

$$R_1 \geq x^* \geq \gamma^* + \lambda > \gamma^* \geq R_1. \quad \square$$

**Remark 7.** Changing slightly the proof of Lemma 6 we could prove that  $i_P(\tilde{F}, B_{R_1}) = 0$  holds also under (H0),(H1),(H5) and (H7). Furthermore, it is worth mentioning that (H7) is equivalent to

$$(H7') \quad \text{there exists } t_0 \in [0, T] \text{ such that } \gamma(t_0) \geq R_1,$$

a condition that is, in general, more readily verifiable.

#### 4. MAIN RESULTS

Now we are ready to formulate and prove our main results.

**Theorem 8.** *If the hypotheses (H0)-(H3), (H5) and (H6) hold, then problem (1.3) has at least one solution  $x$  such that  $c_m R_1 \leq x(t) \leq R_2$  on  $[0, T]$ . The same conclusion remains true with (H3) replaced by (H4) and/or (H6) replaced by (H7).*

*Proof.* Let the operators  $F$  and  $\tilde{F}$  be respectively given by (2.6) and (3.1). Recall that  $F$  and  $\tilde{F}$  coincide on  $\overline{B_{R_2}} \setminus B'_{R_1}$  and thus each fixed point of  $\tilde{F}$  in  $\overline{B_{R_2}} \setminus B'_{R_1}$  is a solution of (1.3).



Now, if  $\tilde{F}x = x$  for some  $x \in \partial B_{R_2} \cup \partial B'_{R_1}$ , then this  $x$  is a solution to our problem. On the other hand, if  $\tilde{F}x \neq x$  for all  $x \in \partial B_{R_2} \cup \partial B'_{R_1}$ , then,  $i_P(\tilde{F}, B_{R_2}) = 1$  by Lemma 5 and  $i_P(\tilde{F}, B'_{R_1}) = 0$  by Lemma 6. Therefore, by Theorem 2 (where we put  $\Omega_P = B_{R_2}$  and  $\Omega'_P = B'_{R_1}$ ), the operator  $\tilde{F}$  has a fixed point in  $B_{R_2} \setminus \overline{B'_{R_1}}$ . To summarize, problem (1.3) has a solution  $x \in \overline{B_{R_2}} \setminus B'_{R_1}$ . Finally, notice that  $x \in \overline{B_{R_2}} \setminus B'_{R_1}$  if and only if  $x \in P$  and  $c_m R_1 \leq x(t) \leq R_2$  on  $[0, T]$ . This completes the proof of the theorem.  $\square$

**Remark 9.** We would like to emphasize that the assumptions of Theorem 8 are weaker than those of Theorem 3.2 of [9]. In particular, if  $f_m(t, x) \leq m^2 R_2$  for  $t \in [0, T]$  and  $x \in [c_m R_2, R_2]$  then (H2) and (H4) of Theorem 8 are satisfied with  $g_1(t) = m^2 R_2$ . Clearly, hypotheses (H2)–(H4) also hold if there exists a continuous function  $\hat{g}$ , such that  $f_m(t, x) \leq \hat{g}(t) R_2$  for  $t \in [0, T]$  and  $x \in [c_m R_2, R_2]$  with

$$\min_{t \in [0, T]} \int_0^T G_m(t, s) \hat{g}(s) \, ds \leq c_m$$

or

$$\max_{t \in [0, T]} \int_0^T G_m(t, s) \hat{g}(s) \, ds \leq 1.$$

Moreover, if  $f_m(t, x) \geq m^2 R_1$  for  $t \in [0, T]$  and  $x \in [c_m R_1, R_1]$  then (H5)–(H7) of Theorem 8 are fulfilled with  $g_0(t) = m^2 R_1$ . For similar type of comparisons see [12, 15].

**Remark 10.** Notice that the nature of our approach does not allow us to derive direct conclusions regarding the uniqueness of the solution in Theorem 8. On the other hand, Theorem 8 provides a localization  $x \in \overline{B_{R_2}} \setminus B'_{R_1}$  of the corresponding solution  $x$  to problem (1.3), which means that

$$(4.1) \quad x \in P \quad \text{and} \quad c_m R_1 \leq x(t) \leq R_2 \quad \text{for } t \in [0, T].$$

Having in mind Remark 7, it is not difficult to modify the proof of Theorem 8 so that, under the assumptions (H0), (H1), (H2), (H3) (or (H4)), (H5) and (H7), we obtain the existence of a solution to problem (1.3) in the set  $\overline{B_{R_2}} \setminus B_{R_1}$ , where in comparison with the localization from Theorem 8, the set  $B'_{R_1}$  is replaced by  $B_{R_1}$ . Observe that this means that the corresponding solution  $x$  of the given problem will satisfy in addition to (4.1) also the condition  $x^* \geq R_1$ .

Next, we will apply Theorem 8 to provide sufficient conditions for the existence of positive solutions of problem (1.2), which is a special case of (1.3) with

$$r(t) = \frac{e(t)}{\mu}, \quad s(t) = \frac{c}{\mu}, \quad \alpha = 1 - 2\mu \quad \text{and} \quad \beta = 1 - \mu.$$

Recall that it was proved in [8] that the necessary condition for the existence of a positive solution of (1.2) is  $\bar{e} > 0$ , and all existence theorems known up to now (cf. [7, 8, 9]) required  $e$  to be strictly positive on  $[0, T]$ . However, it is easy to verify that for

$$(4.2) \quad e(t) = 0.1V_0 + 1.8 + (2.1 - V_0) \cos t - 3 \cos^2 t, \quad T = 2\pi, \quad a = 0, \quad b = 2 \quad \text{and} \quad c = 0.1,$$

with  $V_0 > 3$ , we have  $\bar{e} > 0$  and  $e_* < 0$ , while

$$u(t) = (V_0 - 2) + \cos t,$$

is a solution to the corresponding problem (1.1) (Example (4.2) is in fact, a slight modification of the example by G. Propst from [6, (19)]). This indicates that the positivity of  $e$  can not be a necessary condition for the existence of positive solutions to problem (1.1) and it should be weakened. To our knowledge, next existence principle allows to deal for the first time also with the case  $e_* \leq 0$ .

**Theorem 11.** *Suppose that*

$$(C0) \quad a \geq 0, \quad 0 < \mu < \frac{1}{2}, \quad c > 0, \quad T > 0, \quad \text{and} \quad e : [0, T] \rightarrow \mathbb{R} \text{ is continuous.}$$

Moreover, assume that there are  $m > 0$ ,  $\kappa > 0$ , and  $R_1, R_2 > 0$  with  $R_1 < R_2$  such that (2.4) holds and

$$(C1) \quad e_* \leq 0 \quad \text{and} \quad (c_m R_1)^\mu \geq \frac{c + \sqrt{c^2 - 4\mu m^2 e_*}}{2\mu m^2},$$

$$(C2) \quad (c_m R_1)^{1-2\mu} \geq \kappa \mu,$$

$$(C3) \quad \int_0^T G_m(s, s) e_+(s) \, ds \geq \frac{c_m R_1}{\kappa},$$

$$(C4) \quad e^* \leq c R_2^\mu.$$

Then there exists a positive solution of problem (1.2).

*Proof.* We have  $f_m(t, x) = \frac{e(t)}{\mu} x^{1-2\mu} - \frac{c}{\mu} x^{1-\mu} + m^2 x$ . We will check the hypotheses (H0)–(H2), (H4)–(H6) of Theorem 8. Put  $g_1(t) = m^2 R_2$  and  $g_0(t) = \kappa e_+(t)$  for  $t \in [0, T]$ .

Clearly, (H0) follows from (C0). We will show that (H1) holds, as well. Indeed, (C1) gives  $\mu m^2 x^{2\mu} - c x^\mu + e_* \geq 0$  on  $[c_m R_1, R_2]$ . Hence  $x^{1-2\mu} (\mu m^2 x^{2\mu} - c x^\mu + e_*) \geq 0$  on  $[c_m R_1, R_2]$ . Consequently, we have

$$f_m(t, x) \geq \frac{e_*}{\mu} x^{1-2\mu} - \frac{c}{\mu} x^{1-\mu} + m^2 x \geq 0 \quad \text{for } t \in [0, T] \text{ and } x \in [c_m R_1, R_2].$$

This, together with (2.4) means that (H1) holds.

Now, we will show that (H2) follows from  $\mu \in (0, \frac{1}{2})$ , (C1) and (C4). We need to prove that

$$e^* \leq \mu m^2 R_2 x^{2\mu-1} + c x^\mu - \mu m^2 x^{2\mu}$$

for  $x \in [c_m R_2, R_2]$ . The function  $\mu m^2 R_2 x^{2\mu-1} + c x^\mu - \mu m^2 x^{2\mu}$  is decreasing on  $[c_m R_2, R_2]$ , since by  $\mu \in (0, \frac{1}{2})$  and (C1) we get

$$(\mu m^2 R_2 x^{2\mu-1} + c x^\mu - \mu m^2 x^{2\mu})' = \mu x^{\mu-1} ((2\mu - 1) m^2 R_2 x^{\mu-1} + c - 2\mu m^2 x^\mu) < 0.$$

From (C4) we obtain

$$e^* \leq c R_2^\mu \leq \mu m^2 R_2 x^{2\mu-1} + c x^\mu - \mu m^2 x^{2\mu},$$

which proves (H2). Due to Lemma 1,  $\delta^* = R_2$  in this case. Hence, (H4) is fulfilled, as well.

Let  $t \in [0, T]$  be such that  $e(t) < 0$ . Then, since  $e_+(t) = 0$  in this case, the inequality in (H5) obviously holds. Next, let  $t \in [0, T]$  be such that  $e(t) \geq 0$ . Then we need to show that

$$(4.3) \quad e(t) (x^{1-2\mu} - \kappa \mu) \geq x^{1-\mu} (c - \mu m^2 x^\mu)$$

for  $x \in [c_m R_1, R_1]$ . Since

$$\frac{c + \sqrt{c^2 - 4\mu m^2 e_*}}{2\mu m^2} \geq \frac{c}{\mu m^2},$$

from (C1) and (C2) we get  $c - \mu m^2 x^\mu \leq 0$  and  $x^{1-2\mu} - \kappa \mu \geq 0$  for  $x \in [c_m R_1, R_1]$ . This gives (4.3), and therefore (H5) holds. (H6) follows immediately from (C3).  $\square$

**Remark 12.** Notice that in Theorem 11 the smaller are the difference  $e^* - e_*$  and the period  $T$ , the greater is the chance to find proper  $m, \kappa, R_1$  and  $R_2$ .

Moreover, one can verify that example (4.2) mentioned above does not satisfy the assumptions of Theorem 11. Possible extensions of our existence principle so that they would cover also such examples remain an open problem.

Next example is an illustrative application of Theorem 11.

**Example 13.** Let us consider problem (1.2) with the parameter values  $a = 1.6$ ,  $\mu = 0.01$ ,  $c = 0.005$ ,  $T = 1$  and being  $e$  the continuous periodic function defined on  $[0, 1]$  as follows

$$e(t) = \begin{cases} e_* + \frac{e^* - e_*}{t_1} t & \text{for } t \in [0, t_1), \\ e^* & \text{for } t \in [t_1, t_2), \\ e^* - \frac{e^* - e_*}{1 - t_2} (t - t_2) & \text{for } t \in [t_2, 1], \end{cases}$$

where  $t_1 = 0.0005$ ,  $t_2 = 0.9995$ ,  $e^* = 0.00548239$  and  $e_* = -0.00005$ .

Now, defining  $m = 0.7$ ,  $R_1 = 25$ ,  $R_2 = 10000$  and  $\kappa = 2200$ , direct computations show that all the assumptions of Theorem 11 are satisfied and therefore there exists a positive solution of problem (1.2) (note that, using explicit formulas from Appendix A.1 of [9] and the Mathematica software, we can for  $a = 1.6$  and  $m = 0.7$  approximate the value of  $c_m$  by 0.94144. Similarly we get  $G_m(s, s) \approx 1.96026$ ).

In [9, Example 3.7] we showed that our main result of [9], namely Theorem 3.2, is more general than [8, Theorem 1.8] (see also [9, Corollary 3.6]) when applied to problem (1.2). However, since the function  $e(t) \equiv 1.54$  used in Example 3.7 is constant, problem (1.2) has a constant solution, namely

$$x(t) = \left( \frac{e(t)}{c} \right)^{\frac{1}{\mu}} \equiv \left( \frac{1.54}{1.49} \right)^{100} = 27.1297.$$

Although [9, Example 3.7] is correct, the case of  $e$  constant is not meaningful for the pumping effect in the studied configuration (see [8, Definition 1.2]). Now we present an application of Theorem 8 to problem (1.2) that allows a nonconstant positive function  $e$  which, as it will be shown in Example 15, does not satisfy the assumptions of [8, Theorem 1.8].

**Theorem 14.** *Suppose that (C0) of Theorem 11 holds, and moreover:*

$$(C5) \quad e_* > 0,$$

$$(C6) \quad \text{there is } m > 0 \text{ such that } \frac{c^2}{\mu e_*^2} \left( \frac{1}{c_m^\mu} e^* - e_* \right) \leq m^2 < \left( \frac{\pi}{T} \right)^2 + \left( \frac{a}{2} \right)^2.$$

*Then problem (1.2) has a positive solution  $x$  such that*

$$0 < x(t) \leq \frac{1}{c_m} \left( \frac{e^*}{c} \right)^{\frac{1}{\mu}} \quad \text{for all } t \in [0, T].$$

*Proof.* We will verify that the assumptions of Theorem 8 are satisfied.

*Claim 1.* (H0) and (H1) are true.

Clearly, (H0) is a consequence of (C0). Furthermore, let  $m^2$  be as in (C6) and let

$$R_2 = \frac{1}{c_m} \left( \frac{e^*}{c} \right)^{\frac{1}{\mu}}.$$

We have

$$f_m(t, x) = \frac{e(t)}{\mu} x^{1-2\mu} - \frac{c}{\mu} x^{1-\mu} + m^2 x \quad \text{for } t \in [0, T] \text{ and } x \geq 0.$$

Notice that

$$(4.4) \quad \begin{cases} f_m(t, x) \geq f_*(x) & \text{for all } t \in [0, T] \text{ and } x \geq 0, \text{ where} \\ f_*(x) = \frac{e^*}{\mu} x^{1-2\mu} - \frac{c}{\mu} x^{1-\mu} + m^2 x = \frac{x^{1-2\mu}}{\mu} (e_* - c x^\mu + m^2 \mu x^{2\mu}). \end{cases}$$

Obviously,  $f_*(x) \geq 0$  whenever  $x \in [0, (\frac{e_*}{c})^{\frac{1}{\mu}}]$ . Let  $x \in [(\frac{e_*}{c})^{\frac{1}{\mu}}, R_2]$ . Taking into account (C6), we get

$$\begin{aligned} \mu x^{2\mu-1} f_*(x) &= e_* - c x^\mu + m^2 \mu x^{2\mu} \geq e_* - c R_2^\mu + m^2 \mu \left( \left( \frac{e_*}{c} \right)^{\frac{1}{\mu}} \right)^{2\mu} \\ &= e_* - \frac{e^*}{c_m^\mu} + m^2 \mu \left( \frac{e_*}{c} \right)^2 = \mu \left( \frac{e_*}{c} \right)^2 \left( m^2 - \frac{c^2}{\mu e_*^2} \left( \frac{e^*}{c_m^\mu} - e_* \right) \right) \geq 0. \end{aligned}$$

Thus  $f_*(x) \geq 0$  for all  $x \in [0, R_2]$  and hence

$$f_m(t, x) \geq 0 \quad \text{for } t \in [0, T] \text{ and } x \in [0, R_2].$$

In particular, we conclude that (H1) is true (with an arbitrary  $R_1 \in (0, R_2)$ ).

*Claim 2.* (H2) and (H4) are true.

First, we will prove that  $f_m(t, x) \leq m^2 R_2$  for  $t \in [0, T]$  and  $x \in [c_m R_2, R_2]$ . Indeed, for  $t \in [0, T]$  and  $x \geq 0$  we have

$$f_m(t, x) \leq \frac{e^*}{\mu} x^{1-2\mu} - \frac{c}{\mu} x^{1-\mu} + m^2 x.$$

Furthermore, if  $x \geq c_m R_2 = \left( \frac{e^*}{c} \right)^{\frac{1}{\mu}}$ , then

$$\frac{e^*}{\mu} x^{1-2\mu} - \frac{c}{\mu} x^{1-\mu} = \frac{x^{1-2\mu}}{\mu} (e^* - c x^\mu) \leq \frac{x^{1-2\mu}}{\mu} \left( e^* - c \frac{e^*}{c} \right) = 0.$$

Therefore, for  $t \in [0, T]$  and  $x \in [c_m R_2, R_2]$  we have

$$f_m(t, x) \leq \frac{e^*}{\mu} x^{1-2\mu} - \frac{c}{\mu} x^{1-\mu} + m^2 x \leq m^2 x \leq m^2 R_2.$$

Now, it is easy to show (see also Remark 9) that (H2) and (H4) are true with  $g_1(t) = m^2 R_2$ .

*Claim 3.* (H5) and (H6) are true.

First, we will show that there is  $R_1 \in (0, R_2)$  such that  $f_m(t, x) \geq m^2 R_1$  for  $t \in [0, T]$  and  $x \in [c_m R_1, R_1]$ . Indeed, due to (C5) we can choose  $R_1 \in (0, R_2)$  in such a way that

$$(4.5) \quad (1 - c_m) m^2 \mu R_1^{2\mu} + c R_1^\mu \leq e_* c_m^{1-2\mu}.$$

On the other hand, for  $t \in [0, T]$  and  $x \in [c_m R_1, R_1]$  we deduce

$$\begin{aligned} f_*(x) &= \frac{e_*}{\mu} x^{1-2\mu} - \frac{c}{\mu} x^{1-\mu} + m^2 x \geq \frac{e_*}{\mu} (c_m R_1)^{1-2\mu} - \frac{c}{\mu} R_1^{1-\mu} + m^2 c_m R_1 \\ &= \frac{R_1^{1-2\mu}}{\mu} (e_* c_m^{1-2\mu} - c R_1^\mu + c_m m^2 \mu R_1^{2\mu}) \end{aligned}$$

and hence, by (4.5),

$$f_*(x) \geq m^2 R_1,$$

wherefrom, according to (4.4), the desired inequality immediately follows. Now, it is easy to show (see also Remark 9) that (H5) and (H6) are true with  $g_0(t) = m^2 R_1$ .

By Claims 1–3, all the assumptions of Theorem 8 are satisfied and the proof can be completed applying it.  $\square$


**Example 15.** Let us modify [9, Example 3.7] in order to admit a non-constant function  $e$ . Indeed, consider problem (1.2) with  $a = 1.6$ ,  $\mu = 0.01$ ,  $c = 1.49$ ,  $T = 1$  and  $e$  a continuous periodic function such that  $e_* = 1.54$ . Notice that in this case we cannot apply [8, Theorem 1.8] since

$$\frac{c^2}{4\mu e_*} \approx 36.0406 > 10.5096 \approx \left(\frac{\pi}{T}\right)^2 + \frac{a^2}{4}.$$


On the other hand, for  $m = 0.7$  we have  $c_m \approx 0.9414$  and then Theorem 14 implies that this problem has a positive solution  $x$  provided that  $e^* < 1.5443$ . Moreover,

$$0 < x(t) \leq \frac{1}{c_m} \left(\frac{e^*}{c}\right)^{\frac{1}{\mu}} < 38.0844 \quad \text{for all } t \in [0, T].$$

Finally we show some numerical examples for particular choices of  $e$ :



Solution of problem (1.2) with  $a = 1.6$ ,  
 $\mu = 0.01$ ,  $c = 1.49$ ,  $T = 1$  and  
 $e(t) = 1.54215 + 0.002097 \cos(2\pi t)$



Solution of problem (1.2) with  $a = 1.6$ ,  
 $\mu = 0.01$ ,  $c = 1.49$ ,  $T = 1$  and  
 $e(t) = 1.54215 - 0.02(t - 3t^2 + 2t^3)$

Some computations in Examples 13 and 15 were made with the help of the software system *Mathematica*.

#### ACKNOWLEDGEMENTS

We would like to thank an anonymous referee for his/her careful reading of the manuscript and the many remarks that improved the quality of the paper. In particular his/her insights lead us to example (4.2).

J. A. Cid was partially supported by Ministerio de Educación y Ciencia, Spain, and FEDER, Project MTM2013-43404-P, G. Infante was partially supported by G.N.A.M.P.A. - INdAM (Italy), M. Tvrđý was supported by GA ĀR Grant P201/14-06958S and RVO: 67985840 and M. Zima was partially supported by the Centre for Innovation and Transfer of Natural Science and Engineering Knowledge of University of Rzeszów. This research was completed when M. Zima was visiting Departamento de Matemáticas, Universidade de Vigo, Campus de Ourense, and Mathematical Institute, Czech Academy of Sciences in April, May and June 2016; the kind hospitality is gratefully acknowledged.

#### REFERENCES

- [1] G. Liebau, Über ein ventilloes Pumpprinzip, *Naturwissenschaften* 41 (1954) 327.
- [2] H. Andersson, W. van der Wijngaart, P. Nilsson, P. Enoksson and G. Stemme, A valve-less diffuser micropump for microfluidic analytical systems, *Sensor. Actuat. B-Chem.* 72 (2001) 259–265.
- [3] A. Borzì and G. Propst, Numerical investigation of the Liebau phenomenon, *Z. Angew. Math. Phys.* 54 (2003) 1050–1072.

- [4] T.T. Bringley, S. Childress, N. Vandenberghe and J. Zhang, An experimental investigation and a simple model of a valveless pump, *Phys. Fluids* 20 (2008), 033602, 15 pages.
- [5] J. Männer, A. Wessel and T.M. Yelbuz, How does the tubular embryonic heart work? Looking for the physical mechanism generating unidirectional blood flow in the valveless embryonic heart tube, *Developmental Dynamics* 239 (4) (2010) 1035–1046; <http://onlinelibrary.wiley.com/doi/10.1002/dvdy.22265/full>.
- [6] G. Propst, Pumping effects in models of periodically forced flow configurations, *Phys. D* 217 (2006) 193–201.
- [7] P.J. Torres, *Mathematical models with singularities - A Zoo of Singular Creatures*, Atlantis Briefs in Differential Equations, vol. 1, Atlantis Press, Paris, 2015.
- [8] J.A. Cid, G. Propst and M. Tvrdý, On the pumping effect in a pipe/tank flow configuration with friction, *Phys. D* 273-274 (2014) 28–33.
- [9] J.A. Cid, G. Infante, M. Tvrdý and M. Zima, A topological approach to periodic oscillations related to the Liebau phenomenon, *J. Math. Anal. Appl.* 423 (2015) 1546–1556.
- [10] G. Infante, P. Pietramala and M. Tenuta, Existence and localization of positive solutions for a nonlocal BVP arising in chemical reactor theory, *Commun. Nonlinear Sci. Numer. Simulat.* 19 (2014) 2245–2251.
- [11] K.Q. Lan, Multiple positive solutions of semilinear differential equations with singularities, *J. London. Math. Soc* 63 (2001) 690–704.
- [12] K.Q. Lan, Multiple positive solutions of Hammerstein integral equations and applications to periodic boundary value problems, *Appl. Math. Comput.* 154 (2004) 531–542.
- [13] Y.X. Li, Existence of positive solutions for the cantilever beam equations, *Nonlinear Anal. Real World Appl.* 27 (2016) 221–237.
- [14] J.R.L. Webb, Existence of positive solutions for a thermostat model, *Nonlinear Anal. Real World Appl.* 13 (2012) 923–938.
- [15] J.R.L. Webb, Positive solutions of nonlinear equations via comparison with linear operators, *Discrete Contin. Dyn. Syst.* 33 (2013) 5507–5519.
- [16] P. Omari and M. Trombetta, Remarks on the lower and upper solutions method for second and third-order periodic boundary value problems, *Appl. Math. Comput.* 50 (1992) 1–21.
- [17] M.A. Krasnosel'skiĭ and P.P. Zabreĭko, *Geometrical methods of nonlinear analysis*, Springer-Verlag, Berlin, 1984.
- [18] D. Guo and V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*, Academic Press, Boston, 1988.
- [19] H. Amann, Fixed point equations and nonlinear eigenvalue problems in ordered Banach spaces, *SIAM Rev.* 18 (1976) 620–709.
- [20] K. Deimling, *Nonlinear Functional Analysis*, Springer, New York, 1985.
- [21] E. Zeidler, *Nonlinear Functional Analysis and Its Applications*, Springer, New York, 1993.



JOSÉ ÁNGEL CID, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDADE DE VIGO, 32004, PABELLÓN 3,  
CAMPUS DE OURENSE, SPAIN

*E-mail address:* `angelcid@uvigo.es`

GENNARO INFANTE, DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DELLA CALABRIA,  
87036 ARCAVACATA DI RENDE, COSENZA, ITALY

*E-mail address:* `gennaro.infante@unical.it`

MILAN TVRDÝ, MATHEMATICAL INSTITUTE, CZECH ACADEMY OF SCIENCES, CZ 115 67 PRAHA 1,  
ŽITNÁ 25, CZECH REPUBLIC

*E-mail address:* `tvrdy@math.cas.cz`

MIROSLAWA ZIMA, DEPARTMENT OF DIFFERENTIAL EQUATIONS AND STATISTICS, FACULTY OF MATH-  
EMATICS AND NATURAL SCIENCES, UNIVERSITY OF RZESZÓW, PIGONIA 1, 35-959 RZESZÓW, POLAND.

*E-mail address:* `mzima@ur.edu.pl`